

A generalization of the duality for finite multiple harmonic q -series

Gaku Kawashima

Abstract

Recently, Bradley studied partial sums of multiple q -zeta values and proved a duality result. In this paper, we present a generalization of his result.

Keywords: finite multiple harmonic q -series

1 Introduction

Recently, finite multiple harmonic sums (MHSs for short) have been studied in connection with theoretical physics [1, 10]. In [5, 11], the p -divisibility of MHSs for primes p have been investigated. MHSs have a remarkable property known as the duality and a generalization of this formula, which we call the difference formula for MHSs, was given in [6, Theorem 3.8]. On the other hand, in [2], Bradley proved a q -analog of the duality for MHSs. In the present paper, we shall consider a q -analog of the difference formula for MHSs. We note that the argument is parallel to that in [6].

Here, we explain the duality for finite multiple harmonic q -series due to Bradley. Let $0 < q < 1$. The q -analog of a non-negative integer n is given by

$$[n]_q = \frac{1 - q^n}{1 - q}.$$

For any multi-index (i.e. a finite sequence of positive integers) $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$, we define

$$a_{\boldsymbol{\mu}}(n) = \sum_{n=n_1 \geq \dots \geq n_p \geq 0} \frac{q^{(\mu_1-1)(n_1+1)+\dots+(\mu_p-1)(n_p+1)}}{[n_1+1]_q^{\mu_1} \dots [n_p+1]_q^{\mu_p}}, \quad 0 \leq n \in \mathbb{Z},$$

and

$$b_{\boldsymbol{\mu}}(n) = \sum_{n=n_1 \geq \dots \geq n_p \geq 0} \frac{q^{(n_2+1)+\dots+(n_p+1)}}{[n_1+1]_q^{\mu_1} \dots [n_p+1]_q^{\mu_p}}, \quad 0 \leq n \in \mathbb{Z}.$$

We note that the sum of the infinite series $\sum_{n=0}^{\infty} a_{\boldsymbol{\mu}}(n)$ is the quantity known as the (non-strict) multiple q -zeta value, which has been investigated in recent

years [3, 4, 7, 8, 9]. The following is the duality for finite multiple harmonic q -series:

$$\sum_{i=0}^k (-1)^i q^{\frac{i(i+1)}{2}} \begin{bmatrix} k \\ i \end{bmatrix}_q a_{\boldsymbol{\mu}}(i) = b_{\boldsymbol{\mu}^*}(k), \quad 0 \leq k \in \mathbb{Z}, \quad (1)$$


where

$$\begin{bmatrix} k \\ i \end{bmatrix}_q = \frac{[k]_q!}{[i]_q! [k-i]_q!}$$

is the q -binomial coefficient and $\boldsymbol{\mu}^*$ is the dual multi-index of $\boldsymbol{\mu}$. (The formula is slightly modified from Bradley's one for the purpose of generalization.) The definition of $\boldsymbol{\mu}^*$ will be given in Section 3. For example, we have

$$(2, 2)^* = (1, 2, 1), \quad (1, 1, 2)^* = (3, 1) \quad \text{and} \quad (4)^* = (1, 1, 1, 1) \quad (2)$$

by the diagrams



where the lower arrows are in the complementary slots to the upper arrows.

We next illustrate the main result of this paper. For a multi-index $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$, the quantity $|\boldsymbol{\mu}| = \mu_1 + \dots + \mu_p$ is called the weight of $\boldsymbol{\mu}$. We introduce nested sums

$$c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) = \begin{bmatrix} n+k \\ n \end{bmatrix}_q^{-1} \sum_{\substack{n=n_1 \geq \dots \geq n_p \geq 0 \\ k=k_1 \geq \dots \geq k_r \geq 0}} \frac{q^{(\mu_1-1)(n_1+1) + \dots + (\mu_p-1)(n_p+1) + k_2 + \dots + k_r}}{[n_{i_1} + k_{j_1} + 1]_q \cdots [n_{i_m} + k_{j_m} + 1]_q}, \quad 0 \leq n, k \in \mathbb{Z},$$

for multi-indices $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ and $\boldsymbol{\nu} = (\nu_1, \dots, \nu_r)$ of the same weight m . The subscripts $i_1, \dots, i_m$ and $j_1, \dots, j_m$ are defined by

$$(i_1, \dots, i_m) = (\underbrace{1, \dots, 1}_{\mu_1}, \underbrace{2, \dots, 2}_{\mu_2}, \dots, \underbrace{p, \dots, p}_{\mu_p})$$

and

$$(j_1, \dots, j_m) = (\underbrace{1, \dots, 1}_{\nu_1}, \underbrace{2, \dots, 2}_{\nu_2}, \dots, \underbrace{r, \dots, r}_{\nu_r}),$$

respectively. For example, for $\boldsymbol{\mu} = (3, 1)$ and $\boldsymbol{\nu} = (1, 1, 2)$, we have

$$c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) = \begin{bmatrix} n+k \\ n \end{bmatrix}_q^{-1} \times \sum_{\substack{n=n_1 \geq n_2 \geq 0 \\ k=k_1 \geq k_2 \geq k_3 \geq 0}} \frac{q^{2(n_1+1) + k_2 + k_3}}{[n_1 + k_1 + 1]_q [n_1 + k_2 + 1]_q [n_1 + k_3 + 1]_q [n_2 + k_3 + 1]_q}.$$

The following is the main result of this paper: For any multi-index μ , we have

$$\sum_{i=0}^k (-1)^i q^{\frac{i(i+1)}{2}} \begin{bmatrix} k \\ i \end{bmatrix}_q a_\mu(n+i) = c_{\mu, \mu^*}(n, k), \quad 0 \leq n, k \in \mathbb{Z}. \quad (3)$$

As we see in Section 3, the equality $c_{\mu, \mu^*}(0, k) = b_{\mu^*}(k)$ holds. Hence the formula (3) is a generalization of the formula (1). In Section 2, we interpret the left-hand side of (3) as the k -th q -difference of the sequence $a_\mu \in \mathbb{C}^{\mathbb{N}}$. The proof of (3) is given in Section 3.

2 q -differences of a sequence

In this section, we define the k -th q -difference of a sequence for a non-negative integer k and give an explicit expression for it. Throughout this paper, we fix a complex number q equal to neither 0 nor 1. (When dealing with multiple q -zeta values, we usually assume that $0 < q < 1$. But it is not necessary in finite expressions to restrict q to the range $0 < q < 1$.) In the following, we denote by $\mathbb{N}$ the set of non-negative integers.

Definition 2.1. For any $z \in \mathbb{C}$, we define the difference operator $\Delta_z: \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$ by putting

$$(\Delta_z a)(n) = a(n) - za(n+1)$$

for any $a \in \mathbb{C}^{\mathbb{N}}$ and any $n \in \mathbb{N}$.

Definition 2.2. For any $k \in \mathbb{N}$, we define the k -th q -difference operator by

$$\Delta_{q,k} = \Delta_{q^k} \circ \Delta_{q^{k-1}} \circ \cdots \circ \Delta_q,$$

where $\Delta_{q,0}$ is defined to be the identity on $\mathbb{C}^{\mathbb{N}}$.

Definition 2.3. We define the operator $\nabla_q: \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$ by putting

$$(\nabla_q a)(n) = (\Delta_{q,n} a)(0)$$

for any $a \in \mathbb{C}^{\mathbb{N}}$ and any $n \in \mathbb{N}$.

Let $\mathbb{C}[[X]]$ (resp. $\mathbb{C}[[X, Y]]$) be the ring of formal power series in one variable (resp. two variables) over $\mathbb{C}$. For a sequence $a \in \mathbb{C}^{\mathbb{N}}$, we consider a formal power series

$$F_a(X, Y) = \sum_{n,k=0}^{\infty} (\Delta_{q,k} a)(n) \frac{X^n Y^k}{[n]_q! [k]_q!} \in \mathbb{C}[[X, Y]]. \quad (4)$$

The quantities

$$[n]_q = \frac{1 - q^n}{1 - q} \quad \text{and} \quad [n]_q! = [n]_q [n-1]_q \cdots [1]_q$$

are the q -integer and the q -factorial, respectively. As usual, we put $[0]_q! = 1$. The q -derivative of a formal power series $f(X) \in \mathbb{C}[[X]]$ is defined as

$$D_q f(X) = \left(\frac{d}{dX} \right)_q f(X) = \frac{f(qX) - f(X)}{qX - X} \in \mathbb{C}[[X]].$$

We have the q -Leibniz rule

$$D_q^n (f(X)g(X)) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (D_q^k f)(X) (D_q^{n-k} g)(q^k X) \quad (5)$$

for any $f(X), g(X) \in \mathbb{C}[[X]]$ and any $n \in \mathbb{N}$, where

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}$$

is the q -binomial coefficient. We put $\partial_X = (\partial/\partial X)_q$ and $\partial_Y = (\partial/\partial Y)_q$. For any $f(X, Y) \in \mathbb{C}[[X, Y]]$, we define

$$\Lambda_X f(X, Y) = f(qX, Y) \quad \text{and} \quad \Lambda_Y f(X, Y) = f(X, qY).$$

By the definition of the q -derivative, we have

$$(1-q)X\partial_X = 1 - \Lambda_X \quad \text{and} \quad (1-q)Y\partial_Y = 1 - \Lambda_Y. \quad (6)$$

The q -commutator of operators A and B is defined as

$$[A, B]_q = AB - qBA.$$

We have the following q -commutation relations:

$$\begin{aligned} [\partial_X, \Lambda_X]_q &= [\partial_Y, \Lambda_Y]_q = 0, \\ [\Lambda_X, X]_q &= [\Lambda_Y, Y]_q = 0, \\ [\partial_X, X]_q &= [\partial_Y, Y]_q = 1. \end{aligned} \quad (7)$$

We note that for a formal power series

$$f(X, Y) = \sum_{n,k=0}^{\infty} a(n, k) \frac{X^n Y^k}{[n]_q! [k]_q!} \in \mathbb{C}[[X, Y]]$$

the equality

$$\begin{aligned} & (q\partial_X \Lambda_Y + \partial_Y - 1)f(X, Y) \\ &= \sum_{n,k=0}^{\infty} \{q^{k+1}a(n+1, k) + a(n, k+1) - a(n, k)\} \frac{X^n Y^k}{[n]_q! [k]_q!} \end{aligned} \quad (8)$$

holds. From this, we easily see that

$$(q\partial_X \Lambda_Y + \partial_Y - 1)F_a(X, Y) = 0. \quad (9)$$

Lemma 2.4. *If a formal power series $f(X, Y) \in \mathbb{C}[[X, Y]]$ satisfies two conditions*

$$(q\partial_X \Lambda_Y + \partial_Y - 1)f(X, Y) = 0 \quad \text{and} \quad f(X, 0) = 0,$$

then we have $f(X, Y) = 0$.

Proof. Let

$$f(X, Y) = \sum_{n, k=0}^{\infty} a(n, k) \frac{X^n Y^k}{[n]_q! [k]_q!} \in \mathbb{C}[[X, Y]]$$

satisfy the two conditions of the lemma. Then, by (8), we have

$$q^{k+1}a(n+1, k) + a(n, k+1) - a(n, k) = 0 \quad \text{for any } n, k \in \mathbb{N}$$

and

$$a(n, 0) = 0 \quad \text{for any } n \in \mathbb{N}.$$

Therefore we obtain the result by using induction on k . □

For any sequence $a \in \mathbb{C}^{\mathbb{N}}$, we put

$$f_a(X, Y) = \sum_{n=0}^{\infty} a(n) \frac{(X - qY)(X - q^2Y) \cdots (X - q^nY)}{[n]_q!} \in \mathbb{C}[[X, Y]].$$

We note that

$$\begin{aligned} \partial_X \{ (X - q^mY)(X - q^{m+1}Y) \cdots (X - q^nY) \} \\ = [n - m + 1]_q (X - q^mY)(X - q^{m+1}Y) \cdots (X - q^{n-1}Y) \end{aligned} \quad (10)$$

and

$$\begin{aligned} \partial_Y \{ (X - q^mY)(X - q^{m+1}Y) \cdots (X - q^nY) \} \\ = -q^m [n - m + 1]_q (X - q^{m+1}Y)(X - q^{m+2}Y) \cdots (X - q^nY) \end{aligned} \quad (11)$$

for any integers $1 \leq m \leq n$, which are immediate from the definition of the q -derivative. A q -analog of the exponential function is given by

$$e(X) = \sum_{n=0}^{\infty} \frac{X^n}{[n]_q!} \in \mathbb{C}[[X]].$$

Proposition 2.5. *For any sequence $a \in \mathbb{C}^{\mathbb{N}}$, we have*

$$F_a(X, Y) = f_a(X, Y)e(Y).$$

Proof. It is easily seen that

$$F_a(X, 0) = f_a(X, 0)e(0).$$

According to Lemma 2.4 and (9), we only have to prove the identity

$$(q\partial_X\Lambda_Y + \partial_Y - 1) \{f_a(X, Y)e(Y)\} = 0. \quad (12)$$

By (10), (11) and the q -Leibniz rule (5), we have

$$q\partial_X\Lambda_Y \{f_a(X, Y)e(Y)\} = q \left\{ \sum_{n=1}^{\infty} a(n) \frac{(X - q^2Y) \cdots (X - q^nY)}{[n-1]_q!} \right\} e(qY)$$

and

$$\begin{aligned} & \partial_Y \{f_a(X, Y)e(Y)\} \\ &= -q \left\{ \sum_{n=1}^{\infty} a(n) \frac{(X - q^2Y) \cdots (X - q^nY)}{[n-1]_q!} \right\} e(qY) + f_a(X, Y)e(Y). \end{aligned}$$

From these, the identity (12) immediately follows. $\square$

Corollary 2.6. *Let $a \in \mathbb{C}^{\mathbb{N}}$ be a sequence. Then, for any $n, k \in \mathbb{N}$, we have*

$$(\Delta_{q,k}a)(n) = \sum_{i=0}^k (-1)^i q^{\frac{i(i+1)}{2}} \begin{bmatrix} k \\ i \end{bmatrix}_q a(n+i).$$

Proof. We apply the operator $\partial_X^n \partial_Y^k$ to both sides of the equation in Proposition 2.5:

$$\partial_X^n \partial_Y^k F_a(X, Y) = \partial_X^n \partial_Y^k \{f_a(X, Y)e(Y)\}. \quad (13)$$

The right-hand side is equal to

$$\sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q (\partial_X^n \partial_Y^i f_a)(X, Y) e(q^i Y)$$

by the q -Leibniz rule (5). Since we have

$$(\partial_X^n \partial_Y^i f_a)(0, 0) = (-1)^i q^{\frac{i(i+1)}{2}} a(n+i),$$

the desired equality follows from (13) on setting $X = Y = 0$. $\square$

Corollary 2.7. *Let $a \in \mathbb{C}^{\mathbb{N}}$ be a sequence. Then, for any $n \in \mathbb{N}$, we have*

$$(\nabla_q a)(n) = \sum_{k=0}^n (-1)^k q^{\frac{k(k+1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q a(k).$$

Proof. It follows immediately from Corollary 2.6 on setting $n = 0$. $\square$

3 The difference formula for finite multiple harmonic q -series

We begin with the definition of the dual of a multi-index. A multi-index is a finite sequence of positive integers. For a multi-index $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$, the quantities $|\boldsymbol{\mu}| = \mu_1 + \dots + \mu_p$ and $l(\boldsymbol{\mu}) = p$ are called the weight of $\boldsymbol{\mu}$ and the length of $\boldsymbol{\mu}$, respectively. The multi-indices of weight m are in one-to-one correspondence with the subsets of the set $\{1, 2, \dots, m-1\}$ by the mapping

$$\mathcal{S}_m: (\mu_1, \dots, \mu_p) \mapsto \{\mu_1, \mu_1 + \mu_2, \dots, \mu_1 + \mu_2 + \dots + \mu_{p-1}\}.$$

For example, in the case $m = 3$, we have

$$(3) \mapsto \emptyset, \quad (1, 2) \mapsto \{1\}, \quad (2, 1) \mapsto \{2\} \quad \text{and} \quad (1, 1, 1) \mapsto \{1, 2\}$$

from the diagrams

$$\begin{array}{c} \circ \quad \circ \quad \circ \\ 1 \quad 2 \end{array}, \quad \begin{array}{c} \downarrow \\ \circ \quad \circ \quad \circ \\ 1 \quad 2 \end{array}, \quad \begin{array}{c} \circ \quad \circ \quad \circ \\ 1 \quad 2 \end{array} \quad \text{and} \quad \begin{array}{c} \downarrow \quad \downarrow \\ \circ \quad \circ \quad \circ \\ 1 \quad 2 \end{array}.$$

Definition 3.1. Let m be a positive integer and $\boldsymbol{\mu}$ a multi-index of weight m . Then, we define the dual of $\boldsymbol{\mu}$ by

$$\boldsymbol{\mu}^* = \mathcal{S}_m^{-1}(\mathcal{S}_m(\boldsymbol{\mu})^c),$$

where $\mathcal{S}_m(\boldsymbol{\mu})^c$ denotes the complement of $\mathcal{S}_m(\boldsymbol{\mu})$ in the set $\{1, 2, \dots, m-1\}$.

Examples are given in (2). We note that the equality

$$(l(\boldsymbol{\mu}) - 1) + (l(\boldsymbol{\mu}^*) - 1) = |\boldsymbol{\mu}| - 1 \tag{14}$$

holds for any multi-index $\boldsymbol{\mu}$. Now, we state the definition of the finite multiple harmonic q -series which are considered in this paper.

Definition 3.2. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ be a multi-index. Then, we put

$$a_{\boldsymbol{\mu}}(n) = \sum_{n=n_1 \geq \dots \geq n_p \geq 0} \frac{q^{(\mu_1-1)(n_1+1) + \dots + (\mu_p-1)(n_p+1)}}{[n_1+1]_q^{\mu_1} \dots [n_p+1]_q^{\mu_p}}$$

and

$$b_{\boldsymbol{\mu}}(n) = \sum_{n=n_1 \geq \dots \geq n_p \geq 0} \frac{q^{(n_2+1) + \dots + (n_p+1)}}{[n_1+1]_q^{\mu_1} \dots [n_p+1]_q^{\mu_p}}$$

for any non-negative integer n .

Definition 3.3. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ and $\boldsymbol{\nu} = (\nu_1, \dots, \nu_r)$ be multi-indices of the same weight m . Then, we put

$$c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) = \left[\begin{matrix} n+k \\ n \end{matrix} \right]_q^{-1} \sum_{\substack{n=n_1 \geq \dots \geq n_p \geq 0 \\ k=k_1 \geq \dots \geq k_r \geq 0}} \frac{q^{(\mu_1-1)(n_1+1)+\dots+(\mu_p-1)(n_p+1)+k_2+\dots+k_r}}{[n_{i_1} + k_{j_1} + 1]_q \cdots [n_{i_m} + k_{j_m} + 1]_q}$$

for any non-negative integers n and k , where the subscripts $i_1, \dots, i_m, j_1, \dots, j_m$ are defined by

$$(i_1, \dots, i_m) = (\underbrace{1, \dots, 1}_{\mu_1}, \underbrace{2, \dots, 2}_{\mu_2}, \dots, \underbrace{p, \dots, p}_{\mu_p})$$

and

$$(j_1, \dots, j_m) = (\underbrace{1, \dots, 1}_{\nu_1}, \underbrace{2, \dots, 2}_{\nu_2}, \dots, \underbrace{r, \dots, r}_{\nu_r}).$$

Let $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ be multi-indices of the same weight. Then, it is easily seen that

$$c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, 0) = a_{\boldsymbol{\mu}}(n) \quad (15)$$

for any $n \in \mathbb{N}$. Moreover, by (14), we have

$$c_{\boldsymbol{\mu}, \boldsymbol{\mu}^*}(0, k) = b_{\boldsymbol{\mu}^*}(k) \quad (16)$$

for any $k \in \mathbb{N}$. For any multi-index $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ with $|\boldsymbol{\mu}| \geq 2$, we define a multi-index ${}^-\boldsymbol{\mu}$ by

$${}^-\boldsymbol{\mu} = \begin{cases} (\mu_1 - 1, \mu_2, \dots, \mu_p) & \text{if } \mu_1 \geq 2 \\ (\mu_2, \dots, \mu_p) & \text{if } \mu_1 = 1. \end{cases}$$

We note that

$${}^-(\boldsymbol{\mu}^*) = ({}^-\boldsymbol{\mu})^*. \quad (17)$$

The following lemma states inductive relations of $c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k)$.

Proposition 3.4. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$, $\boldsymbol{\nu} = (\nu_1, \dots, \nu_r)$ be multi-indices of the same weight greater than 1 and n, k non-negative integers.

(i) If $\mu_1 \geq 2$ and $\nu_1 = 1$, then we have

$$q^{-n-k-1} \{[n+k+1]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) - [k]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k-1)\} = c_{{}^-\boldsymbol{\mu}, {}^-\boldsymbol{\nu}}(n, k).$$

(ii) If $\mu_1 = 1$ and $\nu_1 \geq 2$, then we have

$$[n+k+1]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) - [n]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n-1, k) = c_{{}^-\boldsymbol{\mu}, {}^-\boldsymbol{\nu}}(n, k).$$

Proof. Since the proof of (ii) is similar to that of (i), we prove only (i). We have

$$\begin{aligned} & [n+k+1]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) \\ &= \left[\begin{matrix} n+k \\ n \end{matrix} \right]_q^{-1} \sum_{\substack{n=n_1 \geq \dots \geq n_p \geq 0 \\ k \geq k_2 \geq \dots \geq k_r \geq 0}} \frac{q^{(\mu_1-1)(n_1+1)+\dots+(\mu_p-1)(n_p+1)+k_2+\dots+k_r}}{[n_{i_2}+k_{j_2}+1]_q \cdots [n_{i_m}+k_{j_m}+1]_q} \end{aligned}$$

and

$$\begin{aligned} & [k]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k-1) \\ &= [k]_q \frac{[n]_q! [k-1]_q!}{[n+k-1]_q!} \sum_{\substack{n=n_1 \geq \dots \geq n_p \geq 0 \\ k-1=k_1 \geq \dots \geq k_r \geq 0}} \frac{q^{(\mu_1-1)(n_1+1)+\dots+(\mu_p-1)(n_p+1)+k_2+\dots+k_r}}{[n_{i_1}+k_{j_1}+1]_q \cdots [n_{i_m}+k_{j_m}+1]_q} \\ &= \left[\begin{matrix} n+k \\ n \end{matrix} \right]_q^{-1} \sum_{\substack{n=n_1 \geq \dots \geq n_p \geq 0 \\ k-1 \geq k_2 \geq \dots \geq k_r \geq 0}} \frac{q^{(\mu_1-1)(n_1+1)+\dots+(\mu_p-1)(n_p+1)+k_2+\dots+k_r}}{[n_{i_2}+k_{j_2}+1]_q \cdots [n_{i_m}+k_{j_m}+1]_q}. \end{aligned}$$

Therefore we obtain

$$\begin{aligned} & [n+k+1]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) - [k]_q c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k-1) \\ &= \left[\begin{matrix} n+k \\ n \end{matrix} \right]_q^{-1} \sum_{\substack{n=n_1 \geq \dots \geq n_p \geq 0 \\ k=k_2 \geq \dots \geq k_r \geq 0}} \frac{q^{(\mu_1-1)(n_1+1)+\dots+(\mu_p-1)(n_p+1)+k_2+\dots+k_r}}{[n_{i_2}+k_{j_2}+1]_q \cdots [n_{i_m}+k_{j_m}+1]_q}, \end{aligned}$$

from which the result follows immediately. $\square$

We restate Proposition 3.4 in terms of generating functions. For multi-indices $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ of the same weight, we define

$$G_{\boldsymbol{\mu}, \boldsymbol{\nu}}(X, Y) = \sum_{n, k=0}^{\infty} c_{\boldsymbol{\mu}, \boldsymbol{\nu}}(n, k) \frac{X^n Y^k}{[n]_q! [k]_q!}.$$

Proposition 3.5. *Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ and $\boldsymbol{\nu} = (\nu_1, \dots, \nu_r)$ be multi-indices of the same weight greater than 1.*

(i) *If $\mu_1 \geq 2$ and $\nu_1 = 1$, then we have*

$$q^{-1} \Lambda_X^{-1} \Lambda_Y^{-1} \left(\frac{1 - q \Lambda_X \Lambda_Y}{1 - q} - Y \right) G_{\boldsymbol{\mu}, \boldsymbol{\nu}}(X, Y) = G_{-\boldsymbol{\mu}, -\boldsymbol{\nu}}(X, Y).$$

(ii) *If $\mu_1 = 1$ and $\nu_1 \geq 2$, then we have*

$$\left(\frac{1 - q \Lambda_X \Lambda_Y}{1 - q} - X \right) G_{\boldsymbol{\mu}, \boldsymbol{\nu}}(X, Y) = G_{-\boldsymbol{\mu}, -\boldsymbol{\nu}}(X, Y).$$

Proof. These are immediate from Proposition 3.4. $\square$

We use Proposition 3.5 in order to prove Theorem 3.8 by induction, from which the main result follows easily. We need two lemmas.

Lemma 3.6. (i) *We have*

$$\begin{aligned} (q\partial_X\Lambda_Y + \partial_Y - 1)q^{-1}\Lambda_X^{-1}\Lambda_Y^{-1}\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right) \\ = q^{-2}\Lambda_X^{-1}\Lambda_Y^{-1}\left(\frac{1-q^2\Lambda_X\Lambda_Y}{1-q} - qY\right)(q\partial_X\Lambda_Y + \partial_Y - 1). \end{aligned}$$

(ii) *We have*

$$(q\partial_X\Lambda_Y + \partial_Y - 1)\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - X\right) = \left(\frac{1-q^2\Lambda_X\Lambda_Y}{1-q} - X\right)(q\partial_X\Lambda_Y + \partial_Y - 1).$$

Proof. (i) By q -commutation relations (7), we have

$$\begin{aligned} [q\partial_X\Lambda_Y + \partial_Y - 1, \frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y]_q \\ = [q\partial_X\Lambda_Y + \partial_Y, \frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y]_q - (1-q)\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right) \\ = (q\partial_X\Lambda_Y + \partial_Y - 1) - (1-q)\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right). \end{aligned}$$

We transpose the second term of the right-hand side to the left-hand side to obtain

$$\begin{aligned} (q\partial_X\Lambda_Y + \partial_Y - q)\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right) - q\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right)(q\partial_X\Lambda_Y + \partial_Y - 1) \\ = q\partial_X\Lambda_Y + \partial_Y - 1. \end{aligned}$$

Multiplying by the operator $q^{-2}\Lambda_X^{-1}\Lambda_Y^{-1}$ from the left, we see that

$$\begin{aligned} (q\partial_X\Lambda_Y + \partial_Y - 1)q^{-1}\Lambda_X^{-1}\Lambda_Y^{-1}\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right) \\ - q^{-1}\Lambda_X^{-1}\Lambda_Y^{-1}\left(\frac{1-q\Lambda_X\Lambda_Y}{1-q} - Y\right)(q\partial_X\Lambda_Y + \partial_Y - 1) \\ = q^{-2}\Lambda_X^{-1}\Lambda_Y^{-1}(q\partial_X\Lambda_Y + \partial_Y - 1), \end{aligned}$$

where we have used the identities

$$\Lambda_X^{-1}\partial_X = q\partial_X\Lambda_X^{-1} \quad \text{and} \quad \Lambda_Y^{-1}\partial_Y = q\partial_Y\Lambda_Y^{-1}.$$

If we transpose the second term of the left-hand side to the right-hand side, we obtain the result.

(ii) By a similar computation as in (i), we obtain

$$\begin{aligned} & [q\partial_X\Lambda_Y + \partial_Y - 1, \frac{1 - q\Lambda_X\Lambda_Y}{1 - q} - X]_q \\ &= (q\partial_X\Lambda_Y + \partial_Y - 1) - q(1 - \Lambda_X)\Lambda_Y - (1 - q)X(\partial_Y - 1) \\ &= (1 - (1 - q)X)(q\partial_X\Lambda_Y + \partial_Y - 1). \end{aligned}$$

The second equality is due to (6). From this, the desired identity is easily derived. $\square$

Lemma 3.7. *The mappings*

$$q^{-2}\Lambda_X^{-1}\Lambda_Y^{-1}\left(\frac{1 - q^2\Lambda_X\Lambda_Y}{1 - q} - qY\right) \quad \text{and} \quad \frac{1 - q^2\Lambda_X\Lambda_Y}{1 - q} - X$$

from $\mathbb{C}[[X, Y]]$ to itself are injections.

Proof. We prove only the first one. The second is similar. Since the mapping $q^{-2}\Lambda_X^{-1}\Lambda_Y^{-1}$ is an injection, we only have to show that the mapping

$$\frac{1 - q^2\Lambda_X\Lambda_Y}{1 - q} - qY$$

is an injection. This is obviously a linear mapping. We suppose that the formal power series

$$f(X, Y) = \sum_{n, k=0}^{\infty} a(n, k) \frac{X^n Y^k}{[n]_q! [k]_q!} \in \mathbb{C}[[X, Y]]$$

is in the kernel of the above operator. Then we have

$$[n + k + 2]_q a(n, k) - q[k]_q a(n, k - 1) = 0$$

for any $n, k \in \mathbb{N}$. By induction on k , we see that $a(n, k) = 0$ for any $n, k \in \mathbb{N}$. This completes the proof. $\square$

Theorem 3.8. *For any multi-index μ , we have*

$$(q\partial_X\Lambda_Y + \partial_Y - 1)G_{\mu, \mu^*}(X, Y) = 0.$$

Proof. The proof is by induction on $|\mu|$. In the case $|\mu| = 1$ (i.e. $\mu = (1)$), the theorem follows directly from

$$G_{(1), (1)}(X, Y) = \sum_{n, k=0}^{\infty} \frac{X^n Y^k}{[n + k + 1]_q!}.$$

Let $\mu = (\mu_1, \dots, \mu_p)$ be a multi-index with $|\mu| \geq 2$. We put $\mu^* = (\mu_1^*, \dots, \mu_r^*)$. If $\mu_1 \geq 2$, noting $\mu_1^* = 1$, we find that

$$q^{-2}\Lambda_X^{-1}\Lambda_Y^{-1}\left(\frac{1 - q^2\Lambda_X\Lambda_Y}{1 - q} - qY\right)(q\partial_X\Lambda_Y + \partial_Y - 1)G_{\mu, \mu^*}(X, Y) = 0$$

from Lemma 3.6 (i), Proposition 3.5 (i), (17) and the hypothesis of induction. According to Lemma 3.7, we have

$$(q\partial_X\Lambda_Y + \partial_Y - 1)G_{\boldsymbol{\mu},\boldsymbol{\mu}^*}(X,Y) = 0.$$

Also in the case $\mu_1 = 1$, we can argue in the same way. Therefore we have completed the proof. $\square$

The following is the main result of this paper.

Corollary 3.9. *Let $\boldsymbol{\mu}$ be a multi-index. Then we have*

$$(\Delta_{q,k}a_{\boldsymbol{\mu}})(n) = c_{\boldsymbol{\mu},\boldsymbol{\mu}^*}(n,k)$$

for any $n, k \in \mathbb{N}$.

Proof. By (15), we have $F_{a_{\boldsymbol{\mu}}}(X,0) = G_{\boldsymbol{\mu},\boldsymbol{\mu}^*}(X,0)$. (The formal power series $F_a(X,Y)$ is defined in (4) for any sequence $a \in \mathbb{C}^{\mathbb{N}}$.) Therefore we obtain

$$F_{a_{\boldsymbol{\mu}}}(X,Y) = G_{\boldsymbol{\mu},\boldsymbol{\mu}^*}(X,Y)$$

from Lemma 2.4, (9) and Theorem 3.8. This implies the corollary. $\square$

As a corollary of Corollary 3.9, we obtain the duality for finite multiple harmonic q -series due to Bradley.

Corollary 3.10. *For any multi-index $\boldsymbol{\mu}$, we have*

$$\nabla_q a_{\boldsymbol{\mu}} = b_{\boldsymbol{\mu}^*}.$$

Proof. Since we have (16), the corollary follows from Corollary 3.9 on setting $n = 0$. $\square$

References

- [1] J. Blümlein and S. Kurth, Harmonic sums and Mellin transforms up to two-loop order, Phys. Rev. D **60** (1999), 014018.
- [2] D. M. Bradley, Duality for finite multiple harmonic q -series, Disc. Math. **300** (2005), 44-56.
- [3] D. M. Bradley, Multiple q -zeta values, J. Algebra. **283** (2005), 752-798.
- [4] D. M. Bradley, On the sum formula for multiple q -zeta values, Rocky Mountain J. Math. **37** (2007), 1427-1434.
- [5] M. Hoffman, Quasi-symmetric functions and mod p multiple harmonic sums, preprint arXiv:math.NT/0401319.
- [6] G. Kawashima, A class of relations among multiple zeta values, J. Number Theory. **129** (2009), 755-788.

- [7] Y. Ohno and J. Okuda, On the sum formula for the q -analogue of non-strict multiple zeta values, Proc. Amer. Math. Soc. **135** (2007), 3029-3037.
- [8] Y. Ohno, J. Okuda and W. Zudilin, Cyclic q -MZSV sum, preprint MPIM2008-31.
- [9] J. Okuda and Y. Takeyama, On relations for the multiple q -zeta values, Ramanujan J. **14** (2007), 379-387.
- [10] J. A. M. Vermaseren, Harmonic sums, Mellin transforms and integrals, Int. J. Mod. Phys. A **14** (1999), 2037-2076.
- [11] J. Zhao, Wolstenholme type theorem for multiple harmonic sums, Int. J. Number Theory **4** (2008), 73-106.

Graduate School of Mathematics
 Nagoya University
 Chikusa-ku, Nagoya 464-8602, Japan
 E-mail: m02009c@math.nagoya-u.ac.jp